




# Tracking the Post-discontinuation Contraceptive Use Pathways in Rwanda. Second Analysis of 2014-15 Demographic And Health Survey

Victor Ndaruhutse,<sup>1</sup> Felix Sayinzoga,<sup>1</sup> Joel Serucaca,<sup>1</sup> William Winfrey,<sup>2</sup> and  
Anne Pfitzer<sup>2</sup>

Affiliations: <sup>1</sup>Rwanda Biomedical Center, <sup>2</sup>USAID's Maternal and Child Survival  
Program


# Family planning discontinuation in Rwanda

Contraceptive Discontinuation  
(% discontinued within 12 months)



# What happens after discontinuation?

Evolution of reproductive status post-discontinuation



# Implications

- Post-discontinuation follow up of clients who don't want to become pregnant should be improved—after the first month, there is almost no use of a new FP method, thus resulting in pregnancies that would be considered unintended.
- Discontinuers of injectable and implants seem to be particularly frustrated—relatively few switch to new methods.
- Correct use of Standard Days Method is a particular problem, given high failure rate.

For more information, please visit  
**[www.mcspprogram.org](http://www.mcspprogram.org)**

This presentation was made possible by the generous support of the American people through the United States Agency for International Development (USAID), under the terms of the Cooperative Agreement AID-OAA-A-14-00028. The contents are the responsibility of the authors and do not necessarily reflect the views of USAID or the United States Government.

[facebook.com/MCSPglobal](https://facebook.com/MCSPglobal)

[twitter.com/MCSPglobal](https://twitter.com/MCSPglobal)